


Remove that Foley Catheter!

Foley Catheters can Cause:


- Infections ↑ 
- Length of Stay ↑
- Cost ↑\$\$
- Patient/Resident Discomfort 😞
- Antibiotic Usage ↑

☀️ Foley Catheters allow patients/residents to stay in bed which in turn increases skin breakdown, DVT's, & pneumonia due to their immobility.

**PREVENTION
IS KEY.**

Remove that Foley Catheter!

Foley Catheters can Cause:

- Infections ↑ 
- Length of Stay ↑
- Cost ↑\$\$
- Patient/Resident Discomfort 😞
- Antibiotic Usage ↑

☀️ Foley Catheters allow patients/residents to stay in bed which in turn increases skin breakdown, DVT's, & pneumonia due to their immobility.

**PREVENTION
IS KEY.**

Remove that Foley Catheter!

Foley Catheters **are** indicated for:

- Urinary Tract Obstruction
- Gross Hematuria with clots
- Neurogenic bladder with retention
- Urologic surgery or studies
- Sacral decubiti Stage 3-4 w/incontinence
- Hospice, Comfort or Palliative Care (if patient/resident requests)

Foley Catheters are **not** indicated for:

- Incontinence
- Immobility
- Obtaining urine specimens
- Close monitoring of outputs (outside ICU's)
- Patient/resident request/convenience

Remove that Foley Catheter!

Foley Catheters **are** indicated for:

- Urinary Tract Obstruction
- Gross Hematuria with clots
- Neurogenic bladder with retention
- Urologic surgery or studies
- Sacral decubiti Stage 3-4 w/incontinence
- Hospice, Comfort or Palliative Care (if patient/resident requests)

Foley Catheters are **not** indicated for:

- Incontinence
- Immobility
- Obtaining urine specimens
- Close monitoring of outputs (outside ICU's)
- Patient/resident request/convenience